



Faculty of Resource Science and Technology

**ISOLATION OF ANTIBIOTIC RESISTANCE AMONG
PSEUDOMONAS AERUGINOSA FROM AQUACULTURE
ENVIRONMENT AND RIVERS IN SARAWAK**

Soh Khar Mun

**Bachelor of Science with Honours
(Resource Biotechnology)
2015**

**QW
131
M963
2015**

Isolation of Antibiotic Resistance among *Pseudomonas aeruginosa* from Aquaculture Environment and Rivers in Sarawak

Soh Khar Mun (38898)

This project is submitted in fulfillment of the requirement for the Degree of Bachelor of
Science with honors
(Resource Biotechnology)

Supervisor: Dr. Samuel Lihan

Co-supervisor: Dr. Mohd Azlan Jayasilan

Resource Biotechnology Program
Department of Molecular Biology

Faculty of Resource Science and Technology
Universiti Malaysia Sarawak

Acknowledgment

I would like to express my gratitude to my supervisor, Dr. Samuel Lihan for spending time to guide and advise me throughout the project. Other than that, I would like to appreciate Miss Leong Sui Sien, Miss Joyce anak Liang, Miss Kathleen Michelle Mikal, Miss Magarate Rita Ak Elvis, and Mr. Toh Seng Chiew, postgraduate students of Virology lab from Department of Molecular Biology. I would also like to acknowledge Mr. Iskandarshah bin Abang Hassimsah, the lab assistant of Virology lab, all resource personnel from University Malaysia Sarawak (UNIMAS) for their patient guidance and concerns throughout my Final Year Project. My special thanks also go to the board of University Malaysia Sarawak (UNIMAS) for providing all facilities, apparatus, and materials for my research.

Also, my further appreciation to Miss Nurul Aqilah Aqiludeen, master students from Faculty of Medicine and Health Science, UNIMAS for her advises during my research. I would also like to acknowledge Brigita Aquaculture at Kampung Haji Baki, Asia Aquaculture (M) Sdn Bhd at Sampadi and Bako for giving me the opportunity to conduct my research. Also, my special thanks go to Mr. Asia, Mr. Abu, and Mr. Michael for their precious time and help throughout my sampling trip.

Next, I would like to express my gratitude and appreciate to my family members especially my parents for their love, patience, continuous financial and emotional support during my stress and depress moment. Last but not least, I would also like to thank my friends and lab-mates, Chai Shin Nei, Cheng Wei Kang, Lee Sai Yeng, and Tang Lok Ing for their caring, support, and sharing their knowledge and ideas with me to solve the problems I encountered.

UNIVERSITI MALAYSIA SARAWAK

Grade: _____

Please tick (✓)

Final Year Project Report

Masters

PhD



DECLARATION OF ORIGINAL WORK

This declaration is made on the 10th day of June 2015.

Student's Declaration:

I SOH KHAR MUN, 38898, FACULTY OF RESOURCE SCIENCE AND TECHNOLOGY
----- (PLEASE INDICATE STUDENT'S NAME, MATRIC NO. AND FACULTY) hereby
declare that the work entitled ISOLATION OF ANTIBIOTIC RESISTANCE PSEUDOMONAS AERUGINOSA FROM
----- AQUACULTURE ENVIRONMENT AND RIVERS IN SARAWAK
is my original work. I have not copied from any other students' work or from any other
sources except where due reference or acknowledgement is made explicitly in the text, nor
has any part been written for me by another person.

10/6/2015

Date submitted

SOH KHAR MUN (38898)

Name of the student (Matric No.)

Supervisor's Declaration:

I DR. SAMUEL LIHAN (SUPERVISOR'S NAME) hereby certifies that
the work entitled Isolation of Antibiotic Resistance of Pseudomonas aeruginosa
by the above named student, and was submitted to the from Aquaculture Environment and Rivers in Sarawak (TITLE) was prepared
fulfillment for the conferment of Bachelor of Science with Honours (Resource Biotechnology) as a * partial/full
(PLEASE INDICATE THE DEGREE), and the aforementioned work, to the best of my
knowledge, is the said student's work.

Received for examination by: DR. SAMUEL LIHAN Date: 10-06-2015
(Name of the supervisor)

I declare that Project/Thesis is classified as (Please tick (✓)):


- ☐ **CONFIDENTIAL** (Contains confidential information under the Official Secret Act 1972)*
☐ **RESTRICTED** (Contains restricted information as specified by the organisation where research was done)*
☒ **OPEN ACCESS**

Validation of Project/Thesis


I therefore duly affirm with free consent and willingly declare that this said Project/Thesis shall be placed officially in the Centre for Academic Information Services with the abiding interest and rights as follows:

- This Project/Thesis is the sole legal property of Universiti Malaysia Sarawak (UNIMAS).
- The Centre for Academic Information Services has the lawful right to make copies for the purpose of academic and research only and not for other purpose.
- The Centre for Academic Information Services has the lawful right to digitalise the content for the Local Content Database.
- The Centre for Academic Information Services has the lawful right to make copies of the Project/Thesis for academic exchange between Higher Learning Institute.
- No dispute or any claim shall arise from the student itself neither third party on this Project/Thesis once it becomes the sole property of UNIMAS.
- This Project/Thesis or any material, data and information related to it shall not be distributed, published or disclosed to any party by the student except with UNIMAS permission.

Student signature

 (10/6/2015)
(Date)

Supervisor signature:


(Date) 10-06-2015

Current Address:

210, LORONG KUANG RANTING, TAMAN KEPONG, 52100 KUALA LUMPUR.

Notes: * If the Project/Thesis is **CONFIDENTIAL** or **RESTRICTED**, please attach together as annexure a letter from the organisation with the period and reasons of confidentiality and restriction.

[The instrument is duly prepared by The Centre for Academic Information Services

Table of Contents

	Page
Acknowledgements	I
Declaration	II
Table of Contents	VI
List of Abbreviation	VII
List of Tables	IX
List of Figures	X
Abstract	1
Abstrak	1
1.0 Introduction	2
2.0 Literature review	4
2.1 Rivers	4
2.2 Aquaculture Industry	4
2.3 <i>Pseudomonas</i>	5
2.3.1 Phylogeny and Taxonomy	5
2.3.2 Morphology	5
2.3.3 Habitat	6
2.3.4 Physiology	6
2.3.5 Epidemiology	7
2.3.6 <i>Pseudomonas aeruginosa</i>	7
2.4 Biochemical Characterization of <i>P. aeruginosa</i>	8
2.5 Antibiotic Resistance of <i>P. aeruginosa</i>	8
3.0 Materials and Methods	10
3.1 Samples Collection	10
3.2 Isolation of <i>P. aeruginosa</i>	11
3.3 Cultural Characterization of <i>P. aeruginosa</i>	12
3.4 Biochemical Characterization of <i>P. aeruginosa</i>	13
3.4.1 MacConkey Agar Test	13
3.4.2 Oxidase Test	13
3.4.3 Triple Sugar Iron (TSI) Agar Test	14
3.4.4 Citrate Utilization Test	14

3.4.5 Catalase Test	14
3.4.6 Starch Hydrolysis Test	15
3.4.7 Nitrate Reduction Test	15
3.5 Molecular Characterization of <i>P. aeruginosa</i>	16
3.5.1 Bacterial Isolates DNA Extraction	16
3.5.1.1 Preparation of Overnight Culture	16
3.5.1.2 DNA Extraction	16
3.5.2 PCR Amplification	17
3.5.3 Agarose Gel Electrophoresis	18
3.5.4 16S rRNA gene Sequencing	18
3.5.4.1 DNA Purification using QIAquick ® Gel Extraction Kit (Qiagen)	18
3.5.4.2 16S rRNA Gene Sequencing	19
3.6 Antibiotic Susceptibility Test	20
3.6.1 Preparation of Bacterial Culture	20
3.6.2 Disk Diffusion Method	20
4.0 Results and Discussions	21
4.1 Samples Collection	21
4.2 Isolation of <i>P. aeruginosa</i>	23
4.3 Cultural Characterization of <i>P. aeruginosa</i>	25
4.4 Biochemical Characterization of <i>P. aeruginosa</i>	29
4.4.1 MacConkey Agar Test	29
4.4.2 Oxidase Test	31
4.4.3 Triple Sugar Iron (TSI) Agar Test	33
4.4.4 Citrate Utilization Test	36
4.4.5 Catalase Test	38
4.4.6 Starch Hydrolysis Test	40
4.4.7 Nitrate Reduction Test	42
4.4.8 Summary on Biochemical Tests	45
4.5 16S rRNA PCR and DNA Sequencing	45
4.6 Antibiotic Susceptibility Test	48
5.0 Conclusion	53
6.0 Recommendation	54

7.0 References	55
8.0 Appendices	60
Appendix 1: Sample Collection at Sampadi, Lundu	60
Appendix 2: Colony Forming Units of Samples	61
Appendix 3: Work Flow Chart	63
Appendix 4: Boiling-centrifugal Method of DNA Extraction	64
Appendix 5: Purification of DNA from Gel using QIAquick ® Gel Extraction Kit (Qiagen)	65
Appendix 6: Lists of Bacterial Isolates and Origin	66
Appendix 7: Summary of Biochemical Tests	69
Appendix 8: DNA Sequencing Results from 1st Base laboratories	71
Appendix 9: Quality control range for <i>P. aeruginosa</i> ATCC 27853	73

List of Abbreviation

μg	Microgram
μm	Micrometer
AmpC	Ampicillin class C
BLAST	Basic Local Alignment Search Tool
bp	Base pair
CFU	Colony Forming Unit
ClO_3^-	Chlorate
CLSI	Clinical and Laboratory Standards Institute
CV-I	Crystal-violet-iodine
ddH ₂ O	Sterile distilled water
dH ₂ O	Distilled water
DNA	Deoxyribonucleic acid
dsDNA	Double-stranded deoxyribonucleic acid
EPS	Exopolysaccharide
EtBr	Ethidium bromide
ETC	Electron Transport Chain
FeS	Iron (II) sulphate
GDP	Gross Domestic Product
H ₂ O ₂	Hydrogen peroxide
H ₂ S	Hydrogen sulphide
ICUs	Intensive Care Units
KIA	Klingler's Iron Agar
LB	Luria-Bertani Broth

mA	Milli ampere
MgCl ₂	Magnesium chloride
mM	Milli mole
N ₂	Nitrogen gas
N ₂ O	Nitrous oxide
NB	Nutrient Broth
NCBI	National Center for Biotechnology Information
NH ₃	Ammonia
NO	Nitric oxide
NO ₂	Nitrite
NO ₃	Nitrate
<i>P. aeruginosa</i>	<i>Pseudomonas aeruginosa</i>
PCR	Polymerase Chain Reaction
PIA	<i>Pseudomonas</i> Isolation Agar
PNP	p-Nitrophenol
rpm	Revolutions per minute
spp.	Species
TBE	Tris-borate-EDTA
TNTC	Too numerous to count
TSA	Tryptic Soy Agar
TSI	Triple Sugar Iron
UV	Ultraviolet
V	Volt
w/v	Weight per volume

List of Tables

		Page
Table 2.1	Biochemical and physiological tests of the PNP degrading isolate <i>P. aeruginosa</i> DN1	8
Table 3.1	The nucleotide sequence and expected size of amplicon for PCR primers	17
Table 3.2	The volume of PCR master mix for PCR amplifications	17
Table 3.3	Cycling conditions for PCR amplification	18
Table 4.1	Water and sediment samples collected from Asia Aquaculture, Sampadi	21
Table 4.2	Water and sediment samples collected from Asia Aquaculture, Bako 1 and Bako 2	21
Table 4.3	Water samples and cultured species collected from aquaculture farm, Kampung Haji Baki	22
Table 4.4	Water samples collected from rivers	22
Table 4.5	Summary of isolates on CHROMagar™ <i>Pseudomonas</i>	24
Table 4.6	Summary of bacterial isolates morphology after Gram stain	27
Table 4.7	The results of TSI agar test	34
Table 4.8	Nucleotide sequence of bacteria species in NCBI database that shown highest similarity with nucleotide sequence of the bacterial isolates from aquaculture environment	48
Table 4.9	The diameter of inhibition zone for AS-R10(S)	51
Table 4.10	The diameter of inhibition zone for BK2-OLT(S)	51
Table 8.1	Colony forming unit (CFU) for each sample on TSA	61
Table 8.2	Isolates of bacteria with source of origin and cultured species	66
Table 8.3	Summary table of biochemical tests	69
Table 8.4	Nucleotide sequence from DNA sequencing	71
Table 8.5	The quality control range for <i>P. aeruginosa</i> ATCC 27853	73

List of Figures

		Page
Figure 3.1	Sarawak River Basin	10
Figure 4.1	Isolation of <i>P. aeruginosa</i> was done by spreading samples collected onto CHROMagar™ <i>Pseudomonas</i> .	24
Figure 4.2	The cultural characteristics of <i>P. aeruginosa</i> were determined by Gram staining procedure.	28
Figure 4.3	The color change of MacConkey agar after 24 hours incubation.	30
Figure 4.4	The color changes of filter paper strip after 5 to 10 seconds.	32
Figure 4.5	The changes of TSI agar after incubation of 24 hours.	35
Figure 4.6	The color changes of Simmon's citrate agar.	37
Figure 4.7	The formation of gas bubbles for catalase test.	39
Figure 4.8	Formation of hydrolysis zone for starch hydrolysis test.	41
Figure 4.9	The development of red color for nitrate reduction test after 2 minutes.	44
Figure 4.10	Purified DNA was shown in 1% pre-stained EtBr agarose gel.	47
Figure 4.11	Antibiotic susceptibility test was performed using disk diffusion method.	52
Figure 8.1	Water and sediment collection at Asia Aquaculture (M) Sdn Bhd, Sampadi.	60

Isolation of antibiotic resistance among *Pseudomonas aeruginosa* from aquaculture environment and river in Sarawak

Soh Khar Mun

Resource Biotechnology Program
Faculty of Science and Technology
Universiti Malaysia Sarawak

ABSTRACT

Pseudomonas aeruginosa, a Gram-negative bacillus-shaped bacterium can be found in various environments. The high adaptability of *P. aeruginosa* is devoted by the simple nutritional requirement and thus contributed *P. aeruginosa* as an opportunistic pathogen. In this study, *P. aeruginosa* was isolated from water and fish in rivers and aquaculture farm in Sarawak. A total of thirty-eight isolates were successfully isolated using CHROMagarTM *Pseudomonas*. The isolates were subjected to a series of biochemical tests such as gram staining, MacConkey agar test, oxidase test, triple sugar iron agar test, citrate test, catalase test, starch hydrolysis test, and nitrate reduction test. Based on the biochemical tests, two isolates possess the characteristics of *P. aeruginosa*. These isolates were further confirmed through molecular identification using 16S rRNA PCR analysis and DNA sequencing. These two *P. aeruginosa* were screened against nine antibiotics: piperacillin, ceftazidime, imipenem, meropenem, gentamicin, tobramycin, amikacin, ciprofloxacin, and norfloxacin using the disk diffusion assay. Both isolates were susceptible to all the antimicrobial agents tested. This suggested that all these antimicrobial agents are effective against *P. aeruginosa*.

Key words: *Pseudomonas aeruginosa*, aquaculture, biochemical test, 16S rRNA gene sequencing, antibiotic resistance

ABSTRAK

Pseudomonas aeruginosa, sejenis Gram-negatif bakteria berbentuk basillus boleh didapati dalam pelbagai persekitaran. *P. aeruginosa* mempunyai penyesuaian yang tinggi disebabkan keperluan khasiat yang mudah dan ini telah menyebabkan *P. aeruginosa* sebagai patogen oportunistik. Dalam kajian ini, *P. aeruginosa* telah diambil dari air dan ikan di sungai dan ladang akuakultur di Sarawak. Sebanyak tiga puluh lapan isolat berjaya diasingkan dengan menggunakan CHROMagarTM *Pseudomonas*. Isolat ini diuji dengan satu siri ujian biokimia seperti pewarnaan gram, ujian agar MacConkey, ujian oxidase, ujian agar tiga gula besi, ujian sitrat, ujian katalase, ujian hidrolisis kanji hidrolisis, dan ujian pengurangan nitrat. Berdasarkan ujian biokimia, dua isolat yang memenuhi ciri-ciri *P. aeruginosa*. Isolat ini seterusnya dikenal pasti melalui pengenalan molekul dengan menggunakan analisis 16S rRNA PCR penjujukan DNA. Kedua-dua *P. aeruginosa* ini telah diskinkan terhadap Sembilan antibiotik: piperacillin, ceftazidime, imipenem, meropenem, gentamicin, tobramycin, amikacin, ciprofloxacin, dan norfloxacin dengan menggunakan kaedah penyerapan cakera. Kedua-dua isolat terdedah kepada semua ejen antimikrob yang diuji. Ini mencadangkan bahawa semua ejen antimikrob berkesan untuk menentang *P. aeruginosa*.

Kata kunci: *Pseudomonas aeruginosa*, akuakultur, ujian biokimia, penjujukan gen 16S rRNA, rintangan antibiotik

1.0 Introduction

Aquaculture industry started in Malaysia since 1920 and today, this sector is one of the most important activities that contributed international trade, foreign exchange, and a significant source of animal protein (Hashim, 2008). In order to reach Malaysia's vision to become a high income country by 2020, the development of aquaculture into large-scale commercial industry is necessary. The increase demand for fish products especially in rural areas and the failure of captured fisheries to produce necessary fish products make aquaculture a major focus industry (Ling *et al.*, 2011). Therefore, the quality control of the production of fishery products is essential as consumption of fishery products which are raw or contaminated may poses health risks to the consumers.

Sarawak River, an important river basin in the state consists of two major tributaries. According to Nyambar and Bong (2010), these two principal tributaries are Sarawak River Kiri and Sarawak River Kanan. The total length of Sarawak River is 120 km with a total area of 2,456.04 km². Kuok *et al.* (2011) reported that Sarawak River covers towns like Bau, Batu Kawa, and Siniawan. In Sarawak, the establishment of aquaculture sector into commercial scale is markedly feasible due to the availability of water and land (Ling *et al.*, 2011).

Pseudomonas aeruginosa is one of the fish pathogens and can be found in soil and water. *P. aeruginosa*, a Gram-negative and non-fastidious bacterium with bacillus-shaped is highly adaptable in various environments. The bacterium requires only simple nutrient requirements and hence is an opportunistic pathogen that can cause infections in human, plant, and animals (Czekajlo-Kolodziej *et al.*, 2006; Franzetti & Scarpellini, 2007; Hirakawa *et al.*, 2010). Apart from that, the ability of *P. aeruginosa* as a multidrug resistant (MDR) and extremely drug resistant (XDR) made the treatment of disease a

difficult task since the bacterium capable to develop resistance during treatment course in a rapid manner (Lister *et al.*, 2009).

In this study, *P. aeruginosa* were isolated from aquaculture farm located in Sampadi, Bako, and Kampung Haji Baki. Besides that, cultured catfish from Kampung Haji Baki was also caught to isolate *P. aeruginosa*. In addition, water samples were collected from rivers located in Permai, Sarawak River Kiri (Muara Tuang and Bintawa), and Sarawak River Kanan (Siniawan and Waterfront). The detection of *P. aeruginosa* was performed using CHROMagar™ *Pseudomonas* as *Pseudomonas* spp. appears as blue colonies on the selective agar. Following that, the bacteria were characterized using conventional biochemical tests and identified through 16S rRNA gene sequencing. Antibiotic susceptibility test was then conducted and the pattern of antibiotic resistance pattern was analyzed.

The objectives of this study were:

1. to isolate *P. aeruginosa* from aquaculture environment and rivers
2. to determine the biochemical characteristics of isolated *P. aeruginosa* through bio-typing
3. to identify *P. aeruginosa* strains using 16S rRNA gene sequencing
4. to determine the antibiotic resistance pattern of *P. aeruginosa*

2.0 Literature Review

2.1 Rivers

Sarawak River consists of two major tributaries, is an important river basin in the state. The two principal tributaries are Sarawak River Kiri and Sarawak River Kanan. The total length of Sarawak River is 120 km which covers a total area of 2,456.04 km² (Nyambar & Bong, 2010). According to Kuok *et al.* (2011), Sarawak River is originated from Kapuas Mountains and discharged into South China Sea after collecting stormwater and wastewater from drains and tributaries. Sarawak River plays major role to serve residents in Kuching City for domestic and industrial use, to support tourism sector in the state, and function as tool to mitigate flood (Kuok *et al.*, 2011). Therefore, the rivers need to be preserved and managed properly.

2.2 Aquaculture Industry

In Malaysia, aquaculture industry started since 1920 from polyculturing Chinese carps in ex-mining pools. Today, vast variety of species such as marine shrimps, freshwater species, shellfishes, and marine finfishes are cultured using brackish water aquaculture, freshwater pond aquaculture, and marine aquaculture (Hashim, 2008). Aquaculture sector is now one of the most important activities which contributed 1.73% to Gross Domestic Product (GDP) in the country. Other than contributing to international trade and foreign exchange, aquaculture industry is also the cheapest source of animal protein especially for the rural communities (Hashim, 2008).

2.3 *Pseudomonas*

2.3.1 Phylogeny and Taxonomy

The genus *Pseudomonas* was first discovered by Carle Gessard from bluish-green pus in 1882 and this genus received names due to its blue-green coloration during culture (Lister *et al.*, 2009). However, the genus was described further in a vague manner in 1894. The name *Pseudomonas* created by Migula came from the combination of Greek and Latin words, pseudo- (false, Greek) and monas (single unit, Latin). Thus, term *Pseudomonas* means false single unit (Siegrist, 2010). *Pseudomonas* is one of the members of Gamma Proteobacteria class of bacteria and belongs to the Pseudomonadaceae family (Moore *et al.*, 2006).

2.3.2 Morphology

Pseudomonas is a Gram-negative bacterium with slightly curved or straight bacillus-shaped (Koneman, 1997; Czekajlo-Kolodziej *et al.*, 2006; Moore *et al.*, 2006; Franzetti & Scarpellini, 2007) measured from 0.5 to 1.0 μm in diameter by 1.5 to 5.0 μm in length (Moore *et al.*, 2006). Moore *et al.* (2006) stated that not all *Pseudomonas* will follow the traditionally described shape. *Pseudomonas putida* is extremely short while others can be unusually long. The motility of *Pseudomonas* is due to the presence of single polar flagellum and this enables *Pseudomonas* to react to chemical stimuli and localize low concentration organic substrates (Moore *et al.*, 2006).

2.3.3 Habitat

Pseudomonas species is highly adaptable in variety environments. Soil, water, plants and animals are common natural habitats of *Pseudomonas* spp. (Haynes, 1951; Franzetti & Scarpellini, 2007). An environment with a temperature range of 4 to 42 °C, pH from 4 to 8 and rich in organic compounds can likely be a habitat for *Pseudomonas* spp. Other than that, the supply of nitrate (NO_3) as the final electron acceptor enables *Pseudomonas* to survive in anaerobic condition (Moore *et al.*, 2006).

2.3.4 Physiology

The nutritional requirements of *Pseudomonas* spp. are simple (Franzetti & Scarpellini, 2007), thus the bacterium is also known as non-fastidious organism. Although the metabolism is carried out through respiration, some *Pseudomonas* spp. will carry out arginine fermentation as well as pyruvate fermentation to yield energy (Moore *et al.*, 2006). Other than that, some *Pseudomonas* species utilize NO_3 as final electron acceptor in nitrate respiration, a supplementary metabolic pathway. In Moore *et al.* (2006), *Pseudomonas chloritidismutans* utilized chlorate (ClO_3^-) as a substitute of energy generating electron acceptor. On the other hand, *Pseudomonas* spp. can use variety of organic compounds as the source of carbon and energy in order to survive in the environment that lacks nutrients (Igbinosa *et al.*, 2012).

2.3.5 Epidemiology

Pseudomonas is an opportunistic pathogen that causes infections in humans, animals and plants (Czekajlo-Kolodziej *et al.*, 2006; Franzetti & Scarpellini, 2007; Hirakawa *et al.*, 2010) due to ubiquitous lifestyle and nutritional diversity. *Pseudomonas* spp. can also be found in hospital, intensive care unit and can be isolated from hand creams, bar soaps, germicide solutions for cleaning, and antiseptic solutions (Bruun *et al.*, 1976).

2.3.6 *Pseudomonas aeruginosa*

Among all *Pseudomonas* spp., *P. aeruginosa* is the most common species to be studied for the pathogenicity. *P. aeruginosa* is an opportunistic pathogen which leads immunocompromised patients to diseases like neutropenia, hematologic cancers, pneumonia, and sickle cell anemia (Igbinosa *et al.*, 2012). On the other hand, *P. aeruginosa* also results in chronic pulmonary disease for patients with cystic fibrosis. According to Shigeki *et al.* (2011), *P. aeruginosa* can produce many virulence factors such as exotoxin and biofilm and causes the bacterium to be a virulent pathogen.

In the research carried out by Stehling *et al.* (2008, 2010), the pathogenesis of *P. aeruginosa* is due to the production of extracellular enzymes and these enzymes, or known as the exozymes include alkaline protease, elastase, exotoxin A, exoenzyme S and hemolysin. These exozymes are produced through clinical infection and involved in the development of animal infections since they will break down host membrane and lyse cell.

2.4 Biochemical Characterization of *P. aeruginosa*

In Ningthoujam and Shovarani (2008), morphological, biochemical, and physiological tests were performed in order to characterize *P. aeruginosa* strain DN1 degrading p-nitrophenol. These include visualization of morphology under light microscope, motility test by hanging drop method, catalase test, oxidase test using oxidase disc, casein and starch hydrolysis, citrate utilization, Indole, MR, VP, NR, gelatin liquefaction, urease production, and Ulrich milk peptonization. Pigment production was also determined using different media such as LB, *Pseudomonas* F medium, *Pseudomonas* P medium, and PIA. The results were shown in Table 2.1.

Table 2.1: Biochemical and physiological tests of the PNP degrading isolate *P. aeruginosa* DN1

Tests	Results
Catalase	+
Casein hydrolysis	+
Starch hydrolysis	-
MacConkey's agar	+(growth as nonfermenter)
EMB (Eosine Methylene Blue) agar	+
Citrate	+
Motility	+
Glucose fermentation semi solid medium	-
Lactose broth	-
Indole	-
MR	-
VP	-
Gelatin liquefaction	+(rapid liquefaction)
Urea broth	+
NR	+
Ulrich Milk broth	Deep red(peptonization)
Oxidase	+
Lipase	+(showing clear halo)

(Source: <http://www.scialert.net/fulltext/?doi=jm.2008.345.351&org=10>)

2.5 Antibiotic Resistance of *P. aeruginosa*

In recent years, the emergence of *P. aeruginosa* strains which were resistant to multidrug was observed worldwide. The ability of *P. aeruginosa* to resist to multiple antimicrobial agents made this bacterium an opportunistic pathogen. The bacterium does not affect

healthy human. In fact, the bacterium causes alternation to the host defense systems (Gales *et al.*, 2001).

Resistance of antipseudomonal β -lactams and new generation cephalosporins, monobactams, and carbapenems were well described. In 1991, *P. aeruginosa* that produced Metallo-Beta-Lactamase (MBL) was first reported from Japan (Manoharan *et al.*, 2010). The resistance to β -lactams is due to several mechanisms such as AmpC β -lactamase production, extended-spectrum β -lactamases, a barrier to diffuse at outer membrane, and efflux mechanisms. The formation of bacteria biofilm also contributed to the antibiotic resistance of *P. aeruginosa* (Gales *et al.*, 2001). These mechanisms limited therapeutic options for diseases.

According to the study conducted by Lister *et al.* (2009), *P. aeruginosa* exhibited highest resistant rates to fluoroquinolones like ciprofloxacin and levofloxacin. This trend was reported from patients in Intensive Care Units (ICUs). The patients in ICUs also have higher rate of resistance than hospitalized patients towards β -lactamase (Lister *et al.*, 2009).

3.0 Materials and Methods

3.1 Samples Collection

Samples were collected from rivers and aquaculture farms. Twelve samples collected were from the surface water of Permai River, Sarawak River Kiri (Bintawa industrial zone and Muara Tuang area), and Sarawak River Kanan (Siniawan and Waterfront area). Figure 3.1 shows Sarawak River Basin. Four water samples were collected Permai River while two samples were collected from each of the Waterfront, Muara Tuang, Bintawa, and Siniawan rivers. The water samples were taken using sterile falcon tubes and kept in icebox before transporting to laboratory for processing.




Figure 3.1: Sarawak River Basin. Sarawak River Basin has two major tributaries which are Sarawak River Kiri and Sarawak River Kanan.

Four aquaculture farms located at Sampadi, Bako 1, Bako 2, and Kampung Haji Baki were selected for samples collection. The samples collected from aquaculture farms include water, sediment, and cultured species. All water and sediment samples from Asia Aquaculture farm at Lundu (Appendix 1), Bako 1, Bako 2, and aquaculture farm, Kampung Haji Baki were taken using sterile falcon tubes while *Clarias gariepinus* or catfish (local name: *ikan keli*) was caught freshly using sterile polyethylene bags from Kampung Haji Baki. All samples were then kept in icebox containing ice and transported to laboratory within 24 hours for processing.

Eighteen water and sediment samples were collected from Asia Aquaculture farm located at Sampadi, Lundu. Twenty-seven water samples were collected from Asia Aquaculture farm at Bako, Kuching, sixteen water and sediment samples were collected from Bako 1 while eleven water and sediments samples were collected from Bako 2. For aquaculture farm at Kampung Haji Baki, thirteen water samples and a catfish were collected. All the samples were then labeled properly and processed immediately upon arrival in the laboratory.

3.2 Isolation of *P. aeruginosa*

Homogenization of water and sediment samples was performed by flicking the falcon tubes. Then, 50 μ L of each water and sediment samples collected were taken directly from and spread onto CHROMagar™ *Pseudomonas* using spread plate technique. The surface, mouth, intestine (front, middle, end), and anus of catfish were swabbed using sterile cotton bud and spread onto CHROMagar™ *Pseudomonas* for isolation of *P. aeruginosa*. Meanwhile, the water and sediment samples, surface, mouth, intestine, and anus of catfish

were also spread plated onto TSA to obtain the total bacteria count of bacteria in the samples collected. After incubation at 37 °C for 24 hours, bacterial colonies formed were enumerated. The bacteria present in the samples were expressed as:

$$\text{Colony forming units (CFU)} = \frac{\text{Direct bacteria cell count} \times \text{Dilution factor}}{\text{Volume plated (mL)}}$$

For the isolation of *P. aeruginosa*, one or two single colonies from each CHROMagar™ *Pseudomonas* plate were randomly selected before sub-culturing onto TSA agar slant for further study and storage purposes. The isolated *P. aeruginosa* were further characterized using conventional biochemical tests such as Gram staining, MacConkey agar test, oxidase test, Triple Sugar Iron (TSI) test, citrate utilization test, catalase test, starch hydrolysis test, and nitrate reduction test (refer Appendix 3).

3.3 Cultural Characterization of *P. aeruginosa*

Gram staining method was carried out according to Cappuccino and Sherman (2004). A drop of distilled water was placed on the center of slide before a small amount of bacterial colonies were placed onto the slide. The distilled water was then spread into a thin area and heat fixation was performed.

The heat-fixed smear of cell was flooded with crystal violet for 1 minute and then washed gently with tap water. The smear was flooded with Gram's iodine mordant for a minute before gently washing procedure. Next, the smear was decolorized with 95% ethanol until the ethanol runs almost clear. Then, the smear was washed gently using tap water. Lastly, the smear was counterstained with safranin for 45 seconds and washed gently with tap water.